



α-Amylase measurements are used as an aid to monitor and diagnose **pancreatitis** and other **pancreatic diseases**.

METHOD: DIRECT (CNPG₃); KINETIC

FEATURES:

FOR THE QUANTITATIVE MEASUREMENT OF α-AMYLASE

Genzyme Direct Amylase Reagent

The Genzyme Direct Amylase assay involves the use of a chromogenic substrate, 2-chloro-4-nitrophenol (CNPG₃). The methodology enables direct measurement of α-amylase without the use of auxiliary enzymes. It is intended for the quantitative measurement of α-amylase activity in serum and plasma.

- ✓ One part, stable liquid; ready to use reagent
- ✓ Short reaction time with no lag phase
- ✓ Excellent precision
- ✓ No significant interferences from triglyceride, glucose, ascorbic acid, bilirubin or hemoglobin

BENEFITS:

- ✓ Easy to use, no additional reagent preparation required
- ✓ Faster throughput on automated analyzers
- ✓ Confidence in results
- ✓ High reliability of testing and fewer re-tests

ORDERING INFORMATION

CONFIGURATION

CATALOG NUMBER

- GENZYME DIRECT AMYLASE REAGENT
- GENZYME DIRECT AMYLASE REAGENT

- 1L
- 10L

- 80-3844-03
- 80-3603-01



THE AMERICAS
 31 New York Avenue
 Framingham, MA 01701-9322
 Phone: 800 332 1042
 Fax: 800 762 6311
 Email: usdiagcustomerservice@genzyme.com

INTERNATIONAL
 50 Gibson Drive
 Kings Hill, West Malling
 Kent, ME19 4AF, UK
 Phone: +44 (0) 1732 220022
 Fax: +44 (0) 1732 220024/5
 Email: ukdiagcustomerservice@genzyme.com

