



GGT measurements are used as an aid to monitor and diagnose **liver diseases.**

METHOD: ENZYMATIC (GPNA); KINETIC

FEATURES:

FOR THE QUANTITATIVE MEASUREMENT OF GGT
Genzyme GGT Reagent

Genzyme's GGT (Gamma-Glutamyl Transferase) assay incorporates a carboxylated substrate for detection of GGT, following a modification of the International Federation of Clinical Chemistry (IFCC) rapid kinetic procedure. It is intended for the measurement of GGT in serum or plasma.

- ✓ Two part stable liquid reagent
- ✓ No significant interference displayed from samples with elevated levels of lipemia, icterus or ascorbic acid
- ✓ Extended linear range to 1200 U/L
- ✓ Serum or plasma acceptable

BENEFITS:

- ✓ Easy to use, no additional preparation required
- ✓ Accurate results
- ✓ Reduces the need for dilutions and repeat testing
- ✓ Laboratory flexibility

ORDERING INFORMATION	CONFIGURATION	CATALOG NUMBER
<input type="radio"/> GENZYME GGT	R1 1 X 100 mL R2 1 X 20 mL	334-10
<input type="radio"/> GENZYME GGT	R1 3 X 100 mL R2 1 X 60 mL	334-30
<input type="radio"/> GENZYME DC-CAL CALIBRATOR	5 X 3 mL	SE-035
<input type="radio"/> GENZYME DC-TROL LEVEL 1	10 X 5 mL	SM-052
<input type="radio"/> GENZYME DC-TROL LEVEL 2	10 X 5 mL	SM-056



THE AMERICAS
 31 New York Avenue
 Framingham, MA 01701-9322
 Phone: 800 332 1042
 Fax: 800 762 6311
 Email: usdiagcustomerservice@genzyme.com

INTERNATIONAL
 50 Gibson Drive
 Kings Hill, West Malling
 Kent, ME19 4AF, UK
 Phone: +44 (0) 1732 220022
 Fax: +44 (0) 1732 220024/5
 Email: ukdiagcustomerservice@genzyme.com

