



Uric acid measurements are used as an aid to monitor and diagnose **gout** and **renal disease**.

METHOD: ENZYMATIC (URICASE); ENDPOINT

FEATURES:

FOR THE QUANTITATIVE MEASUREMENT OF URIC ACID

Genzyme Uric Acid-SL Reagent

The Genzyme Uric Acid procedure uses a spectrophotometric method that is highly specific for the measurement of uric acid in serum.

- ✓ One part stable liquid
- ✓ Applicable to multiple chemistry platforms
- ✓ Wide linear range

BENEFITS:

- ✓ Easy to use, no additional reagent preparation required
- ✓ Flexible laboratory testing
- ✓ Reduces repeat testing and the need for sample dilutions

ORDERING INFORMATION	CONFIGURATION	CATALOG NUMBER
<input type="radio"/> GENZYME URIC ACID-SL	R1 2 X 100 mL	237-60
<input type="radio"/> GENZYME DC-CAL CALIBRATOR	5 X 3 mL	SE-035
<input type="radio"/> GENZYME DC-TROL LEVEL 1	10 X 5 mL	SM-052
<input type="radio"/> GENZYME DC-TROL LEVEL 2	10 X 5 mL	SM-056



THE AMERICAS
 31 New York Avenue
 Framingham, MA 01701-9322
 Phone: 800 332 1042
 Fax: 800 762 6311
 Email: usdiagcustomerservice@genzyme.com

INTERNATIONAL
 50 Gibson Drive
 Kings Hill, West Malling
 Kent, ME19 4AF, UK
 Phone: +44 (0) 1732 220022
 Fax: +44 (0) 1732 220024/5
 Email: ukdiagcustomerservice@genzyme.com

